

Order-Parameter Fluctuations in Ising Spin Glasses at Low Temperatures

Matteo Palassini,^{1,*} Marta Sales,² and Felix Ritort³

¹*Laboratoire de Physique Théorique et Modèles Statistiques,
Bât. 100, Université Paris-Sud, Orsay Cedex 91405, France*

²*Department of Chemical Engineering, Northwestern University, 2145 Sheridan Rd., Evanston, IL-60202 (USA)*

³*Departament de Física Fonamental, Facultat de Física,
Universitat de Barcelona, Diagonal 647, 08028 Barcelona (Spain)*

(Dated: August 14, 2003)

We present a numerical study of the order-parameter fluctuations for Ising spin glasses in three and four dimensions at very low temperatures and without an external field. Accurate measurements of two previously introduced parameters, A and G , show that the order parameter is not self-averaging, consistent with a zero-temperature thermal exponent value $\theta' \simeq 0$, and confirm the validity of the relation $G = 1/3$ in the thermodynamic limit in the whole low-temperature phase, as predicted by stochastic stability arguments.

PACS numbers: 75.40.Mg, 75.10.Nr

I. INTRODUCTION

Understanding the low-temperature physics of short-range spin glasses¹ remains a major unsolved problem. Much of the current debate concentrates on the equilibrium thermodynamics of the Edwards-Anderson model with Ising spins (EAI model), the canonical short-range spin glass. Since analytical approaches pose formidable difficulties, the problem is often studied numerically. However, the existence of large barriers between low-energy configurations has limited so far numerical calculations to small systems sizes, from which it is hard to draw definite conclusions on the large-volume limit.

Two main issues have been addressed in many numerical studies: the existence and character of the finite-temperature spin-glass transition, and the nature of the low-temperature spin-glass phase. A central quantity of interest in the description of the spin-glass phase is the scaling exponent θ' governing the typical energy of the lowest-lying excitations with linear size of order l , which is assumed to scale as $E \sim l^{\theta'}$. In general, θ' may be distinct^{2,3} from the stiffness exponent θ measured in domain-wall computations^{4,5,6,7}, and the stability of the spin-glass phase requires to have $\theta' \geq 0$. In a “many-state” picture⁸ such as the replica-symmetry-breaking picture inspired by mean-field theory^{1,9,10}, one has $\theta' = 0$, hence there are excitations whose energy remains finite (of the order of the coupling strength between two spins) even as their length scale diverges. In a “two-state” picture, such as the droplet model^{11,12}, one has $\theta' > 0$, hence the energy of large-scale excitations diverges with their size. In this case the identity $\theta' = \theta$ is often assumed.

Both the spin-glass transition and the ordered phase have been usually investigated numerically by computing sample-averaged quantities such as the Binder cumulant¹³ or the distribution of the order parameter (OP) and related observables. Recently^{14,15}, it was observed that useful information on both issues can be drawn from the sample-to-sample fluctuations of the OP.

In particular, two dimensionless measures of the OP fluctuations were considered^{14,15}: A , the normalized fluctuation of the spin-glass susceptibility, and G , a ratio between two cumulants of the OP distribution. These two parameters are related to the Binder cumulant, B , via the relation $B = 1 - A/(2G)$. For a model without time-reversal symmetry (TRS), A , G , and B are given by Eqs. (2), (3), and (4) below. The parameter G serves as a good indicator of the existence of phase transitions¹⁵ in systems lacking TRS (for which B is generally a bad indicator), as recently shown for several systems, including Migdal-Kadanoff spin glasses¹⁶, RNA folding models¹⁷, chiral spin systems¹⁸, and mean-field models such as the SK model (with and without a magnetic field), the infinite-range p -spin model¹⁹, and the infinite-range Potts model²⁰. The parameter A has also been studied before for random diluted models at criticality^{21,22}.

In this paper, we investigate the OP fluctuations in the EAI model with Gaussian couplings in three and four dimensions by low-temperature Monte Carlo simulations. We study the case with no external field, which satisfies TRS. In three dimensions (3D), numerical data are available in the literature for A in the high-temperature phase²³ and for G near the critical point²⁴, for the “ $\pm J$ ” coupling distribution. In four dimensions (4D), G was measured at moderately low temperatures, also for the $\pm J$ distribution¹⁴. Here, we study much lower temperatures than in these studies, in order to reduce crossover effects associated to the critical point^{25,26}, which complicate the interpretation of the numerical data at higher temperatures.

A summary of our results is as follows. First, we estimate θ' from the system-size dependence of A , finding, for the system sizes we could reach, a small value of θ' incompatible with the accepted values of the domain-wall exponent ($\theta \simeq 0.2$ in 3D^{4,5,7} and $\theta \simeq 0.7$ in 4D⁶), and compatible with zero. This agrees with recent determinations of θ' from ground-state perturbation methods^{2,3,27} and from low-temperature measurements of the OP distribution^{10,26,28,29} (which all consider sample-averaged

quantities), and supports a picture of the spin-glass phase characterized by two distinct exponents, $\theta > 0$ and $\theta' = 0$. The result $\theta' = 0$ implies that the OP is not *self-averaging* in the thermodynamic limit.

Second, we find good evidence that the identity $G = 1/3$ holds in the whole spin-glass phase in the thermodynamic limit, confirming the validity of sum rules proposed by Guerra³⁰ and first derived for the SK model, which follow from the property of “replica equivalence”^{31,32}.

Third, we find that A and G allow to locate the spin-glass transition reasonably well although, as expected due to TRS and as previously numerically observed²⁴, B provides a better determination (a much more accurate determination is provided by the correlation length²⁴, which we do not investigate here).

We do not study in this paper the surface fractal dimension d_s of the excitations, which is the other exponent, besides θ' , characterizing the spin-glass phase (in particular, $d_s = 0$ in the standard replica-symmetry-breaking picture¹⁰, $d_s > 0$ in the droplet model, while the “TNT” picture^{2,3} predicts the “mixed” behavior $d_s > 0$, $\theta' = 0$).

The rest of the paper is organized as follows. In Sec. II we introduce the different models and observables studied, and discuss the theoretical predictions for these observables. In Sec. III we present and analyze our numerical results for the quantities A , G and B . Finally, in Sec. IV we summarize our conclusions.

II. MODELS, OBSERVABLES, AND THEORETICAL PREDICTIONS

We study the EAI model defined by the Hamiltonian

$$\mathcal{H} = - \sum_{\langle i,j \rangle} J_{ij} S_i S_j, \quad S_i = \pm 1, \quad (1)$$

where L^D Ising spins S_i sit on a (hyper-)cubic lattice in D dimensions with linear size L and periodic boundary conditions in all directions. The couplings J_{ij} are drawn from a Gaussian distribution of zero mean and unit variance. We consider two different models: (i) the case with interactions $\langle i, j \rangle$ restricted to nearest neighbors (referred to as NN model) in $D = 3$ and 4; (ii) the case with interactions restricted to nearest, next-nearest, and next-next-nearest neighbors (referred to as NNN model) in $D = 3$, which has a coordination number $z = 26$.

The NN model has been extensively studied and is known to display a finite-temperature continuous spin-glass transition for $D \geq 3$. Recent estimates of the critical temperature for Gaussian-distributed couplings are $T_c = 0.95 \pm 0.04$ in 3D³³ and $T_c = 1.80 \pm 0.03$ in 4D³⁴.

The NNN model has been much less studied. In Ref. 23, the 3D case with $\pm J$ couplings was considered, but no conclusive evidence of a finite-temperature transition was obtained, the data being compatible with both $T_c \approx 3.27$ and a zero-temperature singularity. Incidentally, in the NNN case we do not expect a large difference

in T_c between binary and Gaussian distributions, due to the large coordination number. For the NNN model, we did not consider temperatures as low as for the NN model, but focused on the phase transition region. Another of the results of this paper is a convincing evidence that indeed a finite-temperature transition exists in this model in 3D.

We measure A , G , and B as a function of temperature T and size L using the following definitions:

$$A(T, L) = \frac{\overline{\langle q^2 \rangle^2} - \overline{\langle q^2 \rangle}^2}{\overline{\langle q^2 \rangle}^2} \quad (2)$$

$$G(T, L) = \frac{\overline{\langle q^2 \rangle^2} - \overline{\langle q^2 \rangle}^2}{\overline{\langle q^4 \rangle} - \overline{\langle q^2 \rangle}^2} \quad (3)$$

$$B(T, L) = \frac{1}{2} \left(3 - \frac{\overline{\langle q^4 \rangle}}{\overline{\langle q^2 \rangle}^2} \right) \quad (4)$$

where $q = \frac{1}{L^D} \sum_i S_i^a S_i^b$ is the spin overlap of two independent systems S_i^a and S_i^b with the same random couplings, and $\langle \dots \rangle$ and $\overline{\langle \dots \rangle}$ stand for thermal and disorder averages, respectively. The OP for a given realization of the disorder is $\langle q^2 \rangle$, therefore A is nothing but the normalized sample-to-sample variance of the OP.

In the paramagnetic phase, $T > T_c$, and for sufficiently large L so that $L \gg \xi$ (where ξ is the correlation length), the OP follows a Gaussian distribution and all three parameters vanish as $1/L^D$, as follows from the central limit theorem. Following the terminology of Wiseman and Domany²¹, this means that the OP is *strongly self-averaging*.

At $T = T_c$, the correlation length diverges and the central limit theorem cannot be applied. For strongly disordered systems such as spin glasses it is known that the OP is not self-averaging at criticality^{21,22}, namely A tends to a finite value in the thermodynamic limit. If A is finite then clearly G must be finite, and standard renormalization-group arguments show that B is also finite at T_c . Since B and G are dimensionless and monotonic in T , in plots of these quantities as a function of T , the curves for different values of L must all cross at $T = T_c$, and one can use this to determine T_c . From standard finite-size scaling, one can then determine the critical exponent ν . Since much work has been devoted to measuring ν from the standard observables (see for instance Refs. 24,33,35,36,37), and we are primarily interested in the low-temperature phase here, we will not attempt a precise determination.

In the spin-glass phase, $T < T_c$, A is expected^{38,39} to vanish linearly with T according to the scaling law

$$A(T, L) \sim T L^{-\theta'} \quad (5)$$

where θ' is the exponent discussed in the Introduction. This law holds under two hypothesis (both satisfied in the case of continuous couplings studied here): (i) the ground state is unique; (ii) the probability distribution

of the energy of the lowest-lying excitations has finite weight at zero energy^{38,39}.

From the above scaling law we see that if $\theta' > 0$, then A vanishes for $L \rightarrow \infty$, namely the OP is *weakly self-averaging* (where “weakly” indicates²¹ that the OP fluctuations vanish more slowly than $1/L^d$, a consequence of the inequality $\theta' < d$). This situation is encountered in the droplet model^{11,12}, as discussed in the Introduction, and also in mean-field models with a marginally stable replica-symmetric solution at low temperatures (such as the spherical SK model⁴⁰). If $\theta' = 0$, as in a “many-state” picture, A remains finite in the thermodynamic limit, namely the OP is *not* self-averaging.

Turning now to G , it is known^{31,32} that in the SK model the following relation holds for $T < T_c$:

$$\lim_{L \rightarrow \infty} G(T, L) = 1/3. \quad (6)$$

According to Guerra³⁰, this relation should hold (for $T < T_c$) in any model which is “stochastically stable” with respect to a mean-field perturbation and which has a non-self-averaging OP. Under the hypothesis (i) and (ii) above, the more general conjecture has also been made⁴⁰ that the above relation holds for $T < T_c$ even if the OP is self-averaging. In this case, G would be finite but both the numerator and the denominator in Eq.(3) would vanish, as for example in the Migdal-Kadanoff spin glass (see Bokil et al. in Ref. 16) and the SK spherical model⁴⁰. It has also been explicitly proven³⁸, under the hypothesis (i) and (ii), that one has $G(T = 0, L) = 1/3$ for any L . Note that models in which $G(T = 0, L) \neq 1/3$ in general will not satisfy the conjecture of Ref. 40.

III. NUMERICAL RESULTS

We simulated the various models with the parallel tempering technique⁴¹, which allows to reach significantly lower temperatures than conventional Monte Carlo methods. The parameters of the simulation are given in Table I. Equilibration of the Monte Carlo runs was tested by monitoring all the measured observables on a logarithmic time scale, checking that they had all converged within their statistical errors, and by applying the equilibration test discussed in Ref. 26.

A. Parameter A

In Figures 1, 2, and 3 we show our numerical results for A in the 3D NN, 4D NN, and 3D NNN models, respectively. The vertical lines in Figures 1 and 2 indicate the estimated value of T_c . In all cases, the behavior of A resembles that observed in the SK model (see Refs. 14,18,19). At high temperatures, A decreases with L , approximately as $1/L^D$, showing that the OP is strongly self-averaging in this regime, as expected. Near

Model	L	$T_{\min}$	$T_{\max}$	N_T	N_s	MCS
3D NN	4	0.1	2.0	18	16000	10^5
	6	0.2	2.0	16	6000	10^5
	8	0.2	2.0	16	6600	10^5
	12	0.94	2.0	14	3751	$3 \cdot 10^5$
	16	0.94	2.0	16	587	10^6
4D NN	3	0.2	2.8	12	16000	10^4
	4	0.2	2.8	12	13951	10^5
	5	0.46	2.8	19	1476	$3 \cdot 10^5$
	7	0.995	2.8	29	832	$3 \cdot 10^5$
3D NNN	4	2.0	5.0	16	9005	10^4
	6	2.0	5.0	16	3258	10^4
	8	2.0	5.0	16	3574	$3 \cdot 10^4$
	12	2.8	5.0	12	1751	10^5
	16	3.4	5.0	9	489	10^5

TABLE I: Parameters of the simulations. L is the linear system size, $T_{\min}$ and $T_{\max}$ the smallest and largest temperatures considered, N_T the number of temperatures in the parallel tempering algorithm, N_s the number of independent realizations of the disorder (samples), and MCS the number of Monte Carlo steps per spin and per temperature.

$T = T_c$, there is a maximum whose position shifts towards T_c as L increases, an effect of finite-size corrections. The shift is modest in the 4D NN model but quite noticeable in the 3D NN model, where even for the largest L the position of the maximum is still significantly larger than T_c . In the 3D NNN model, the position of the maximum is also larger than T_c (see discussion in Sec. III C on the value of T_c in this model), but the shift is less pronounced. The *height* of the maximum increases with L in all models, indicating that A attains a finite value in the thermodynamic limit (since it is bounded from above), namely that the OP is not self-averaging at T_c , as expected.

At low temperatures ($T < T_c$), Figures 1, 2, and 3 show that A is approximately linear in T , in agreement with Eq. (5). Most interestingly, the data for different values of L tend to superimpose to each other. In a scenario with $\theta' > 0$, the data should tend to zero for large L in the whole region below T_c . In 3D, we see no decrease at all in the data with increasing L , while a modest decrease is observed in 4D.

To analyze in more detail the size dependence of A at low temperatures, in Figure 4 we plot the ratio A/T as function of L at different temperatures for the three models. The straight lines represent the scaling law Eq.(5) assuming $\theta' = \theta$, and using the estimates of θ from domain-wall calculations, $\theta = 0.2$ in 3D^{4,5,7} and $\theta = 0.7$ in 4D⁶. No estimates of θ are available for the 3D NNN model, so we use that for the NN model (we expect that θ is a universal exponent equal for both models). Clearly, the data in Figure 4 do not agree with the hypothesis $\theta' = \theta$ for the range of sizes considered, and seem to saturate




FIG. 1: Parameter A for the 3D NN model as a function of the temperature, for different system sizes L . The vertical line represents the estimated value of the critical temperature, $T_c = 0.95 \pm 0.04$.




FIG. 2: Same as Figure 1 but for the 4D NN model. The vertical line corresponds to $T_c = 1.80 \pm 0.03$.

to a constant value instead. We fitted the data with the form $A(T, L)/T = aL^{-\hat{\theta}'}$, where $\hat{\theta}'$ should be seen as an “effective” exponent which depends on the temperature and which effectively takes into account corrections to the leading scaling behavior, with $\hat{\theta}' \rightarrow \theta'$ in the limit $TL^{-\theta'} \rightarrow 0$. The fits give $\hat{\theta}'$ varying from 0.03 ± 0.02 ($T = 0.7$) to 0.00 ± 0.06 ($T = 0.2$) in the 3D NN model, from 0.30 ± 0.05 ($T = 1.0$) to 0.003 ± 0.006 ($T = 0.32$) in the 4D NN model, and from 0.03 ± 0.04 ($T = 2.8$) to 0.08 ± 0.04 ($T = 2.0$) in the 3D NNN model.

Therefore, in all cases the data is compatible with $\theta' = 0$, in agreement with the “many-state” picture, and




FIG. 3: Same as Figure 1 but for the 3D NNN model.

is statistically incompatible with $\theta' = \theta$, in disagreement with the “two-state” picture. As usual, we cannot exclude a crossover²⁵ to a larger value of θ' for larger L . In this case, in the large-volume limit A would be zero at all temperatures, except at $T = T_c$.

A value of θ' compatible with zero was also obtained from the OP distribution^{10,26,28,29,36} and from direct measurements of the energy of low-lying excitations created by perturbing the ground state^{2,3,27}.




FIG. 4: Log-log plot of A/T versus L at different temperatures for the 3D NN model (top), the 4D NN model (middle), and the 3D NNN model (bottom). The straight lines of slope -0.2 (top and bottom) and -0.7 (middle) are the expected scaling behavior of Eq.(5) if $\theta' = \theta$.

B. Parameters G and B

Figures 5, 6 and 7 show our numerical results for G and B for the three models. At high temperatures, G and B vanish approximately as $1/L^D$ in all three models, again indicating strong self-averaging. Near $T = 0$, the data for the NN model in both 3D and 4D are compatible with $G(T = 0, L) = 1/3$ and $B(T = 0, L) = 1$, as expected^{38,40} for a continuous coupling distribution. More importantly, $G(T, L)$ seems to converge to the value $1/3$ for large L in the *whole* low-temperature region, in agreement with Eq.(6). This is particularly evident in the 4D NN model (Figure 6), where $G(T, L)$ has already converged to $1/3$ for $L = 5$ at temperatures below $T \approx T_c/2 \approx 0.9$ (the data points above $1/3$ are due to incomplete equilibration¹⁹). It is quite unlikely that the saturation is a finite-size effect, therefore our results strongly suggest that indeed $G = 1/3$ in the whole spin-glass phase. As discussed above, if A remains finite as $L \rightarrow \infty$ (as indicated by our data), this is an expected consequence of stochastic stability³⁰. If A vanishes, instead, our results would support the more general conjecture of Ref. 40.

C. Critical region

In this subsection we comment on the behavior of G and B near the critical temperature, starting from the NN model.

In 3D, the vertical lines in Figure 5 indicates the position of the critical temperature, using the value $T_c = 0.95 \pm 0.04$ quoted in Ref. 33, which was obtained from the parameter B measured in a large-scale simulation. One sees that the data for both B and G for different values of L come together as T approaches T_c from above, as indicative of a phase transition. Below T_c , the data for G separate again in a statistically significant way, while for B one would need a substantially larger statistics (or larger sizes) to see a clear separation, as observed in previous studies^{33,37}. For example, at $T = 0.82$ the separation between the $L = 12$ and the $L = 4$ data is 1.4 standard deviations for B and 2.5 standard deviations for G . The small separation below T_c is probably due to the vicinity of $D = 3$ to the lower critical dimension^{33,35,37}. We also note that the crossing point of G is at temperatures larger than T_c , and close inspection shows that it shifts towards T_c from above as L increases. A similar shift was observed for the position of the maximum of A in Figure 1.

In 4D, both B and G display a very clear crossing (see Figure 6), as also observed in previous studies^{29,34}. From B we estimate $T_c = 1.80 \pm 0.03$ in agreement with the results of Refs. 34,42. This value is indicated by the vertical lines in Figure 6. As in 3D, the crossing point of G is at temperatures larger than T_c and shifts towards T_c as L increases.

Overall, this confirms that both in 3D and 4D the corrections to scaling are significantly larger for G and A

than for B . Since G and A have also much larger statistical errors than B , the latter quantity is to be preferred to G and A for locating T_c in models with TRS. As already mentioned, a much more accurate quantity for this purpose is the correlation length, which shows a very clear crossing in 3D²⁴, unlike B and G .

Finally, in the 3D NNN model both G and B show a rather clear crossing (see Figure 7). This provides a clear evidence for the existence of a phase transition in 3D Ising spin glasses, confirming recent results for the NN model^{24,35,37} that obtained a convincing evidence (especially Ref.24) after the issue had remained unsolved for a long time. The crossing point is at $T \simeq 3.3$ for B and at somewhat higher temperatures for G , although also here the crossing for G shifts to the left as L increases. From the data for B one might be tempted to conclude that the critical temperature is $T_c \simeq 3.3$. However, if this was the case, the value of B at T_c (which is a universal quantity) would be lower in the 3D NNN model than in the 3D NN model, violating universality. This suggests that the actual value of T_c is significantly lower than 3.3, despite the clear crossing of B (which would then be strongly affected by scaling corrections), and for this reason we have not indicated the position of T_c in Figures 3 and 7. A more detailed analysis⁴³ clearly shows that indeed T_c is significantly lower than 3.3 in this model.




FIG. 5: Parameters G (main figure) and B (inset) for the 3D NN model, as a function of the temperature and for various system sizes. The vertical lines correspond to $T_c = 0.95 \pm 0.04$, the horizontal line in the main figure corresponds to the limit relation $G = 1/3$.

IV. CONCLUSIONS

To conclude, we have provided evidence that the order parameter is not self-averaging in the low-temperature phase of the Edwards-Anderson Ising spin glass in 3D




FIG. 6: Same as Figure 5 but for the 4D NN model. The vertical lines correspond to $T_c = 1.80 \pm 0.03$.




FIG. 7: Same as Figure 5 but for the 3D NNN model. The true critical temperature is significantly lower than the crossing point of B .

and 4D, which implies an exponent $\theta' \simeq 0$, in agreement with a “many-state” picture of the spin-glass phase, such as the replica-symmetry-breaking picture or the “TNT” picture. As usual, due to the limited system sizes that are currently reachable in numerical simulations, we cannot exclude that for larger sizes one recovers self-averaging, nevertheless our result is consistent with other studies which used sample-averaged quantities^{2,3,26,27} and also found $\theta' \simeq 0$. Independently of whether there is self-averaging or not, we have provided evidence that the identity $G = 1/3$ holds in the thermodynamic limit in the whole spin-glass phase, a fact that calls for a theoretical explanation in terms of the geometry and energetics of the low-lying excitations. We have confirmed that G and A can be used to locate the spin-glass transition, although in models with time-reversal symmetry the usual sample-averaged parameters provide a better determination. Finally, we have confirmed the existence of a spin-glass phase transition at finite temperature in three dimensions.

Acknowledgments

We acknowledge support from Spanish Ministerio de Ciencia y Tecnología, Grants BFM2001-3525 (F.R.), AP98-36523875 (M.S.), Generalitat de Catalunya (F.R.). M.P. acknowledges support from the European Commission, contract HPRN-CT-2002-00319 (STIPCO).

* Electronic address: matteo@ipno.in2p3.fr

- ¹ M. Mézard, G. Parisi and M.A. Virasoro, *Spin Glass Theory and Beyond*. World Scientific (Singapore, 1987); K. Binder and A. P. Young, *Rev. Mod. Phys.* **58**, 801 (1986).
- ² F. Krzakala and O.C. Martin, *Phys. Rev. Lett.* **85** 3013 (2000).
- ³ M. Palassini and A.P. Young, *Phys. Rev. Lett.* **85**, 3017 (2000).
- ⁴ A.J. Bray and M.A. Moore, *J. Phys. C* **17**, L463 (1984); W.L. McMillan, *Phys. Rev. B* **29**, 4026 (1984); *Phys. Rev. B* **30**, 476 (1984); M. Cieplak and J. R. Banavar, *J. Phys. A* **23**, 4385 (1990).

- ⁵ A. K. Hartmann, *Phys. Rev. E* **59**, 84 (1999); **60**, 5135 (1999).
- ⁶ A. K. Hartmann, *Phys. Rev. B* **64**, 224430 (1999); K. Hukushima, *Phys. Rev. E* **60**, 3606 (1999).
- ⁷ M. Palassini and A.P. Young, *Phys. Rev. Lett.* **83**, 5126 (1999).
- ⁸ We use the term “state” loosely, namely in the sense of region of the configuration space separated from other regions by high free-energy barriers, and not in the sense of thermodynamic state, for which there is no accepted rigorous definition for spin glasses.
- ⁹ G. Parisi, *Phys. Lett.* **73 A**, 203 (1979); *Phys. Rev. Lett.*

- 43**, 1754 (1979).
- ¹⁰ E. Marinari et al., J. Stat. Phys. **98**, 973 (2000).
 - ¹¹ D. S. Fisher and D. A. Huse, J. Phys. A. **20**, L997 (1987); D. A. Huse and D. S. Fisher, J. Phys. A. **20**, L1005 (1987); D. S. Fisher and D. A. Huse, Phys. Rev. B **38**, 386 (1988).
 - ¹² A. J. Bray and M. A. Moore, in *Heidelberg Colloquium on Glassy Dynamics and Optimization*, L. Van Hemmen and I. Morgenstern eds. (Springer-Verlag, Heidelberg, 1986).
 - ¹³ K. Binder, Z. Phys. B **43**, 119 (1981).
 - ¹⁴ E. Marinari et al., Phys. Rev. Lett. **81**, 1698 (1998).
 - ¹⁵ G. Parisi, M. Picco, and F. Ritort, Phys. Rev. E **60**, 58 (1999).
 - ¹⁶ H. Bokil et al., Phys. Rev. Lett. **82**, 5174 (1999); E. Marinari et al., Phys. Rev. Lett. **82**, 5175 (1999).
 - ¹⁷ A. Pagnani, G. Parisi, and F. Ricci-Tersenghi, Phys. Rev. Lett. **84**, 2026 (2000).
 - ¹⁸ H. Hukushima and K. Kawamura, Phys. Rev. E **61**, R1008 (2000).
 - ¹⁹ M. Picco, F. Ritort, and M. Sales, Eur. Phys. J. B **19**, 565 (2001).
 - ²⁰ K. Hukushima and H. Kawamura, Phys. Rev. E. **62**, 3360 (2000).
 - ²¹ S. Wiseman and E. Domany, Phys. Rev. E **52**, 3469 (1995); Phys. Rev. Lett. **81**, 22 (1998).
 - ²² A. Aharony and A. B. Harris, Phys. Rev. Lett. **77**, 3700 (1996).
 - ²³ E. Marinari, G. Parisi, and F. Ritort, J. Phys. A **27**, 2687 (1994).
 - ²⁴ H. G. Ballesteros et al., Phys. Rev. B **62**, 14237 (2000).
 - ²⁵ H. Bokil, B. Drossel, and M. A. Moore, Phys. Rev. Lett. **81**, 4252 (1998); Phys. Rev. B **62**, 946 (2000).
 - ²⁶ H. G. Katzgraber, M. Palassini, and A. P. Young, Phys. Rev. B **63**, 184422 (2001).
 - ²⁷ E. Marinari and G. Parisi, Phys. Rev. Lett. **86**, 3887 (2001).
 - ²⁸ J.D. Reger, N.R. Bhatt, and A.P. Young, Phys. Rev. Lett. **64**, 1859 (1990).
 - ²⁹ E. Marinari and F. Zuliani, J. Phys. A **32**, 7447 (1999).
 - ³⁰ F. Guerra, Int. J. Mod. Phys. B **10**, 1675 (1996).
 - ³¹ A. P. Young, A. J. Bray, and M. A. Moore, J. Phys. C **17**, L149 (1984).
 - ³² M. Mézard et al., J. Physique **45**, 843 (1984).
 - ³³ E. Marinari, G. Parisi, and J. J. Ruiz-Lorenzo, Phys. Rev. B **58**, 14852 (1998).
 - ³⁴ G. Parisi, F. Ricci-Tersenghi, and J. J. Ruiz-Lorenzo, J. Phys. A **29**, 7943 (1996).
 - ³⁵ M. Palassini and S. Caracciolo, Phys. Rev. Lett. **82**, 5128 (1999).
 - ³⁶ E. Marinari, G. Parisi, and J. J. Ruiz-Lorenzo, in *Spin Glasses and Random Fields*, A. P. Young Ed., World Scientific, Singapore 1998.
 - ³⁷ N. Kawashima and A.P. Young, Phys. Rev. B **53**, R484 (1996).
 - ³⁸ F. Ritort and M. Sales, J. Phys. A **34**, L333 (2001).
 - ³⁹ B. Drossel, H. Bokil, and M. A. Moore, Phys. Rev. E **62**, 7690 (2000).
 - ⁴⁰ F. Ritort and M. Sales, J. Phys. A **33**, 6505 (2000).
 - ⁴¹ K. Hukushima and K. Nemoto, J. Phys. Soc. Jpn. **65**, 1604 (1996).
 - ⁴² N. R. Bhatt and A. P. Young, Phys. Rev. B **37**, 5606 (1988).
 - ⁴³ M. Palassini, in preparation.